

Transport properties near the Anderson transition

C. Villagonzalo, R. A. Römer, and M. Schreiber

Institut für Physik, Technische Universität, D-09107 Chemnitz, Germany
villagonzalo@physik.tu-chemnitz.de

Received dd.mm.yyyy, accepted dd.mm.yyyy by xx

Abstract. The electronic transport properties in the presence of a temperature gradient ∇T in disordered systems near the metal-insulator transition (MIT) are considered. The d.c. conductivity σ , the thermoelectric power S , the thermal conductivity K and the Lorenz number L_0 are calculated for the three-dimensional (3D) Anderson model of localization using the Chester-Thellung-Kubo-Greenwood formulation of linear response. We show that σ , S , K and L_0 can be scaled to one-parameter scaling curves with a single scaling parameter $k_B T / |(\mu - E_c)/E_c|$.

Keywords: disordered solids, metal-insulator transitions, thermoelectric conduction

PACS: 61.43.-j, 71.30.+h, 72.15.Cz

1 Introduction

In this paper we shall demonstrate that in the Anderson model of localization near the metal-insulator transition (MIT) the d.c. conductivity σ , the thermoelectric power S , the thermal conductivity K and the Lorenz number L_0 obey scaling.

A fundamental model often used in dealing with transport phenomena of disordered media is given by the Anderson Hamiltonian [1]. Investigations of this model yield that the energy E dependent conductivity $\sigma(E)$ behaves as

$$\sigma(E) = \begin{cases} \sigma_0 \left| 1 - \frac{E}{E_c} \right|^\nu, & |E| < E_c, \\ 0, & |E| \geq E_c, \end{cases} \quad (1)$$

in energy regions near the mobility edge E_c at which the MIT occurs [2]. Here σ_0 is a constant and ν is a universal critical exponent. Using Eq. (1) we have been able to study the temperature T dependence of σ , S , K and L_0 near the MIT [3] using linear response theory [4]. Here we present our observations of their scaling features.

2 Calculating the transport properties

In the presence of a small temperature gradient ∇T in an open circuit, S is the coefficient of proportionality between ∇T and the electric field it induces. The coefficient K relates ∇T to the heat current while the Lorenz number L_0 measures the ratio between K and the product σT . The transport properties are defined in the framework

of linear response theory as

$$\begin{aligned}\sigma &= L_{11}, & S &= \frac{L_{12}}{|e|TL_{11}}, \\ K &= \frac{L_{22}L_{11} - L_{21}L_{12}}{e^2TL_{11}}, \quad \text{and} \quad L_0 &= \frac{L_{22}L_{11} - L_{21}L_{12}}{(k_B TL_{11})^2},\end{aligned}\tag{2}$$

where e is the electron charge and k_B is Boltzmann's constant. For a noninteracting system with no inelastic scattering processes present, such as phonon mediated hopping, the kinetic coefficients L_{ij} are given in the Chester-Thellung-Kubo-Greenwood formulation [4] as

$$L_{ij} = (-1)^{i+j} \int_{-\infty}^{\infty} A(E) [E - \mu(T)]^{i+j-2} \left[-\frac{\partial f(E, \mu, T)}{\partial E} \right] dE, \tag{3}$$

where $i, j = 1, 2$, μ is the chemical potential of the system, $f(E, \mu, T)$ is the Fermi distribution function, and $A(E)$ describes the system dependent features. In the Anderson model, one finds $A(E)$ to be equal to the critical behavior (1) of σ near the MIT [5].

Previously, we have numerically determined $\mu(T)$ for the 3D Anderson model from its density of states [3]. Using that result, i.e., $\mu(T) \propto T^2$, it is straightforward to determine the kinetic coefficients (3) and, consequently, the transport properties (2).

In order to compare our results with experiments we have chosen the energy units to be in electron volts. For the same reason, we measure σ in units of $\Omega^{-1}\text{cm}^{-1}$. In this paper we only show the results obtained for disorder $W = 12$ where W is the width of the box distribution of randomly chosen potential energies in the Anderson model of localization. We have obtained similar results for other disorders not too close to the critical disorder $W_c \approx 16.5$ [6, 7] and where there are no large fluctuations in the density of states. For $W = 12$, the corresponding mobility edge is at $E_c = 7.5$ [6]. The value of ν is set to 1.3 in agreement with numerical results [8]. Note that this choice of ν determines the magnitude of the transport properties but does not change their T dependence [3] upon which their scaling is based.

3 Scaling Features

Wegner [9] was the first to show for disordered noninteracting electron systems that σ can be written close to the MIT in a scaling form as

$$\sigma(t, \Omega) = b^{2-d} F(b^{1/\nu} t, b^z \Omega). \tag{4}$$

Here F is a system dependent function of the dimensionless parameter t giving the distance from the critical point, Ω is an external parameter such as the frequency, b is a scaling parameter, d is the dimension and z is the dynamical exponent. For the noninteracting case, $z = d$ [10]. In this paper, the appropriate parameters are $d = 3$, $t = |(\mu - E_c)/E_c|$, $\Omega = T$ and $b = t^{-\nu}$ and Eq. (4) simplifies to

$$\frac{\sigma(t, T)}{t^\nu} = F\left(\frac{T}{t^{\nu z}}\right). \tag{5}$$




Fig. 1 Left: The d.c. conductivity obtained in the metallic regime: $E_F - E_c = -0.01$ eV ($\circ$), -0.007 eV (∇), -0.005 eV ($\square$), -0.003 eV ($\triangle$), -0.001 eV ($\diamond$), in the critical regime: $E_F - E_c = 0$ eV ($\bullet$), and in the insulating regime: $E_F - E_c = 0.001$ eV ($+$), 0.003 eV ($\times$). The thin solid line is the scaling curve onto which all data from the different energy regimes collapses. The thick solid line denotes a high temperature expansion [3, 5] while the dashed line is the Sommerfeld expansion [5, 12] for $E_F - E_c = -0.01$ eV. Right: The scaling plot of the thermoelectric power obtained from the different regions in the same way as for σ .

In Fig. 1 we show that the σ data from the metallic ($|E_F| < E_c$), the insulating ($|E_F| > E_c$) and the critical ($E_F = E_c$) energy regions collapse onto a single scaling curve when plotted as a function of $k_B T / |(\mu - E_c)/E_c|$. In the right hand side of Fig. 1, we illustrate similar scaling of S . Note that there is no pre-factor in the scaling form such as in σ since S becomes independent of T at the MIT as $T \rightarrow 0$. It was pointed out in Ref. [5] that S for high and low T might scale in terms of $k_B T / |E_F - E_c|$. This is approximately valid [3] although the proper scaling variable should be $k_B T / |(\mu - E_c)/E_c|$. The scaling curve for S converges at the value $228.4 \mu\text{V/K}$ as predicted in Ref. [11] for $\nu = 1.3$. In the metallic regime, our result agrees with the Sommerfeld expansion result at low T . We find that S grows infinitely large in the insulating regime. This can be attributed to the decrease in charge carriers which contribute to a current.

The corresponding scaling curves for K and L_0 are shown in Fig. 2. They also scale with respect to $k_B T / |(\mu - E_c)/E_c|$. The normalization factor in K is due to the fact that $K \propto k_B T^{\nu+1}$ as $T \rightarrow 0$ at the critical regime. L_0 does not require a pre-factor since it is independent of T at the MIT [11]. The result for L_0 shows that it approaches $\pi^2/3$ in the metallic regime as expected in the Sommerfeld free electron theory of metals. In the critical regime, L_0 converges to 2.414 which is the value for $\nu = 1.3$ predicted in Ref. [11]. Due to the exponential decay of the derivative of the Fermi function as $T \rightarrow 0$, L_0 approaches $\nu + 1 = 2.3$ in the insulating regime.

To summarize, we achieve scaling for the four transport quantities σ , S , K and L_0 with respect to a single parameter $k_B T / |(\mu - E_c)/E_c|$. This factor stems from the form of L_{ij} and from the fact that the explicit T dependence enters via μ and the Fermi distribution function. Comparing our results from σ in Fig. 1 with Eq. (5) we obtain $\nu z = 1$. With $\nu = 1.3$ this yields a dynamical exponent $z = 0.77$




Fig. 2 Left: Scaling plot of the thermal conductivity obtained from the same energy regions as given in Fig. 1. The thin solid line is the scaling curve onto which all data from the different energy regimes collapses. The thick solid line denotes a high temperature expansion [3, 5] while the dashed line is the Sommerfeld expansion [5, 12] for $E_F - E_c = -0.01$ eV. Right: The scaling of the Lorenz number. The additional symbol (*) denotes data obtained deeper in the insulating regime: $E_F - E_c = 0.01$ eV.

which is to be compared to $z = d = 3$ as expected from scaling arguments [9, 10]. Moreover, our result that $\nu z = 1$ seems to violate the Harris criterion [13] which requires $\nu z > 2$. However, this observation needs to be examined in more detail since the Harris criterion might be altered for our choice of T dependence via the Fermi distribution.

We thank D. Belitz, T. R. Kirkpatrick, A. MacKinnon and T. Vojta for useful discussions. Financial support by the DFG through Sonderforschungsbereich 393 is gratefully acknowledged.

References

- [1] P. W. Anderson, Phys. Rev. **109** (1958) 1492
- [2] B. Kramer and A. MacKinnon, Rep. Prog. Phys. **56** (1993) 1469
- [3] C. Villagonzalo, R. A. Römer, and M. Schreiber, cond-mat/9904362, accepted for publication in Eur. Phys. J. B
- [4] G. V. Chester and A. Thellung, Proc. Phys. Soc. **77** (1961) 1005; R. Kubo, J. Phys. Soc. Japan **12** (1957) 570; D. A. Greenwood, Proc. Phys. Soc. **71** (1958) 585
- [5] U. Sivan and Y. Imry, Phys. Rev. B **33** (1986) 551
- [6] H. Grussbach and M. Schreiber, Phys. Rev. B **51** (1995) 663
- [7] K. Slevin and T. Ohtsuki, Phys. Rev. Lett. **82** (1999) 382
- [8] E. Hofstetter and M. Schreiber, Phys. Rev. Lett. **73**, (1994) 3137; E. Hofstetter and M. Schreiber, Phys. Rev. B **49** (1994) 14726
- [9] F. Wegner, Z. Phys. B **25** (1976) 327
- [10] D. Belitz and T. R. Kirkpatrick, Rev. Mod. Phys. **66** (1994) 261
- [11] J. E. Enderby and A.C. Barnes, Phys. Rev. B **49** (1994) 5062
- [12] C. Castellani, C. Di Castro, M. Grilli, and G. Strinati, Phys. Rev. B **37** (1988) 6663
- [13] A. B. Harris, J. Phys. C **7** (1974) 1671